

Referencias bibliográficas:

- Wu, J.M.; Vaughan, C.P.; Goode, P.S.; Redden, D.T.; Burgio, K.L.; Richter, H.E.; Markland, A.D. Prevalence and trends of symptomatic pelvic floor disorders in U.S. women. *Obstet. Gynecol.* 2014, 123, 141–1488.
- Maclennan, A.H.; Taylor, A.W.; Wilson, D.H.; Wilson, D. The prevalence of pelvic floor disorders and their relationship to gender, age, parity and mode of delivery. *BJOG* 2000, 107, 1460–1470.
- Martin-Martin, S.; Pascual-Fernandez, A.; Alvarez-Colomo, C.; Calvo-Gonzalez, R.; Muñoz-Moreno, M.; Cortiñas-Gonzalez, J.R. Urinary incontinence during pregnancy and postpartum. Associated risk factors and influence of pelvic floor exercises. *Arch. Esp. Urol.* 2014, 67, 323–330.
- Rortveit, G.; Hannestad, Y.S. Association between mode of delivery and pelvic floor dysfunction. *Tidsskr. Nor. Lægeforening* 2014, 134, 1848–1852.
- Lan, C.W.; Tavrow, P. Composite measures of women’s empowerment and their association with maternal mortality in low-income countries. *BMC Pregnancy Childbirth* 2017, 17 (Suppl. 2), 337.
- Kenton, K.; Mueller, E.R. The global burden of female pelvic floor disorders. *BJU Int.* 2006, 98 (Suppl. S1), 6–7.
- Wu, J.M.; Vaughan, C.P.; Goode, P.S.; Redden, D.T.; Burgio, K.L.; Richter, H.E.; Markland, A.D. Prevalence and trends of symptomatic pelvic floor disorders in U.S. women. *Obstet. Gynecol.* 2014, 123, 141–1488.
- Maclennan, A.H.; Taylor, A.W.; Wilson, D.H.; Wilson, D. The prevalence of pelvic floor disorders and their relationship to gender, age, parity and mode of delivery. *BJOG* 2000, 107, 1460–1470.
- Mant J, Painter R, Vessey M. Epidemiology of genital prolapse: observations from the Oxford Family Planning Association Study. *Br J Obstet Gynaecol.* 1997 May; 104(5):579-85.
- Swift S, Woodman P, O’Boyle A, Kahn M, Valley M, Bland D, Wang W, Schaffer J. Pelvic Organ Support Study (POSST): the distribution, clinical definition, and epidemiologic condition of pelvic organ support defects. *Am J Obstet Gynecol.* 2005 Mar; 192(3):795-806.
- Sze EH, Sherard GB 3rd, Dolezal JM. Pregnancy, labor, delivery, and pelvic organ prolapse. *Obstet Gynecol.* 2002 Nov; 100(5 Pt 1):981-6.
- Yeung J, Stecher A, Crisp CC, Mazloomdoost D, Smith B, Kleeman SD, Pauls RN. Incidence of Obstetric Anal Sphincter Injuries After Training to Protect the Perineum. *Female Pelvic Med Reconstr Surg.* 2018 Mar/Apr;24(2):126-129. doi: 10.1097/SPV.0000000000000506. PMID: 29474285.
- Eason E, Labrecque M, Wells G, Feldman P. Preventing perineal trauma during childbirth: a systematic review. *Obstet Gynecol.* 2000 Mar;95(3):464-71. doi: 10.1016/s0029-7844(99)00560-8. PMID: 10711565.
- Waszyński E. Zabieg Kristellera--Expressio fetus, jego geneza i współczesne zastosowanie [Kristeller’s procedure--Expressio fetus, its genesis and contemporary application]. *Ginekol Pol.* 2008 Apr;79(4):297-300. Polish. PMID: 18592869.
- Vääräsmäki M, Raudaskoski T. Pregnancy and delivery after a cesarean section. *Duodecim.* 2017;133(4):345-52. PMID: 29205981.
- Barrett G, Pendry E, Peacock J, Victor C, Thakar R, Manyonda I. Women’s sexual health after childbirth. *BJOG.* 2000 Feb;107(2):186-95. doi: 10.1111/j.1471-0528.2000.tb11689.x. PMID: 10688502.
- McDonald EA, Gartland D, Small R, Brown SJ. Frequency, severity and persistence of postnatal dyspareunia to 18 months post partum: A cohort study. *Midwifery.* 2016 Mar;34:15-20. doi: 10.1016/j.midw.2016.01.012. Epub 2016 Feb 8. PMID: 26971442.
- Rathfisch G, Dikencik BK, Kizilkaya Beji N, Comert N, Tekirdag AI, Kadioglu A. Effects of perineal trauma on postpartum sexual function. *J Adv Nurs.* 2010 Dec;66(12):2640-9. doi: 10.1111/j.1365-2648.2010.05428.x. Epub 2010 Aug 23. PMID: 20735499.
- Signorello LB, Harlow BL, Chekos AK, Repke JT. Postpartum sexual functioning and its relationship to perineal trauma: a retrospective cohort study of primiparous women. *Am J Obstet Gynecol.* 2001 Apr;184(5):881-8; discussion 888-90. doi: 10.1067/mob.2001.113855. PMID: 11303195.
- Beckmann MM, Stock OM. Antenatal perineal massage for reducing perineal trauma. *Cochrane Database Syst Rev.* 2013 Apr 30;(4):CD005123. doi: 10.1002/14651858.CD005123.pub3. PMID: 23633325.
- Chen I, Opiyo N, Tavender E, Mortazhejri S, Rader T, Petkovic J, Yogasingam S, Taljaard M, Agarwal S, Laopaiboon M, Wasiak J, Khunpradit S, Lumbiganon P, Gruen RL, Betran AP. Non-clinical interventions for reducing unnecessary caesarean section. *Cochrane Database Syst Rev.* 2018 Sep 28;9(9):CD005528. doi: 10.1002/14651858.CD005528.pub3. PMID: 30264405; PMCID: PMC6513634.

Barca JA, Bravo C, Pintado-Recarte MP, Asúnsolo Á, Cueto-Hernández I, Ruiz-Labarta J, Buján J, Ortega MA, De León-Luis JA. Pelvic Floor Morbidity Following Vaginal Delivery versus Cesarean Delivery: Systematic Review and Meta-Analysis. *J Clin Med.* 2021 Apr 13;10(8):1652. doi: 10.3390/jcm10081652. PMID: 33924472; PMCID: PMC8070303.

Barrett G. et al. Women's sexual health after childbirth. *BJOG: An International Journal of Obstetrics & Gynaecology*, 2000, vol. 107, no 2, p. 186-195.

McDonald EA., et al. Frequency, severity and persistence of postnatal dyspareunia to 18 months post partum: A cohort study. *Midwifery*, 2016, vol. 34, p. 15-20.

Rathfisch G, et al. Effects of perineal trauma on postpartum sexual function. *Journal of advanced nursing*, 2010, vol. 66, no 12, p. 2640-2649.

Signorello L.B., et al. Postpartum sexual functioning and its relationship to perineal trauma: a retrospective cohort study of primiparous women. *American journal of obstetrics and gynecology*, 2001, vol. 184, no 5, p. 881-890.

Abdool, Z.; Lindeque, B.G.; Dietz, H.P. The impact of childbirth on pelvic floor morphology in primiparous Black South African women: A prospective longitudinal observational study. *Int. Urogynecol. J.* 2018, 29, 369–375. [CrossRef] [PubMed]

Huser, M.; Janku, P.; Hudecek, R.; Zbozinkova, Z.; Bursa, M.; Unzeitig, V.; Ventruba, P. Pelvic floor dysfunction after vaginal and cesarean delivery among singleton primiparas. *Int. J. Gynaecol. Obstet.* 2017, 137, 170–173.

Yohay, D.; Weintraub, A.Y.; Mauer-Perry, N.; Peri, C.; Kafri, R.; Yohay, Z.; Bashiri, A. Prevalence and trends of pelvic floor disorders in late pregnancy and after delivery in a cohort of Israeli women using the PFDI-20. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 2016, 200, 35–39.

Moosdorff-Steinhauser, H.F.A.; Albers-Heitner, P.; Weemhoff, M.; Spaanderman, M.E.A.; Nieman, F.H.M.; Berghmansa, B. Factors influencing postpartum women's willingness to participate in a preventive pelvic floor muscle training program: A web-based survey. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 2015, 195, 182–187.

Lipschuetz, M.; Cohen, S.M.; Liebergall-Wischnitzer, M.; Zbedat, K.; Hochner-Celnikier, D.; Lavy, Y.; Yagel, S. Degree of bother from pelvic floor dysfunction in women one year after first delivery. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 2015, 191, 90–94.